

Corporation obtaining approval, the name of its representative, and the address of its main office

Du Pont Kabushiki Kaisha
 Minoru Amoh, President
 2-11-1, Nagata-chou, Chiyoda-ku,
 Tokyo

Approved Type 1 Use Regulation

| | |
|--|---|
| Name of the type of Living Modified Organism: | Maize resistant to Lepidoptera and tolerant to glufosinate herbicide (Modified <i>vip3A</i> , <i>cry2A.127</i> , <i>cry1A.88</i> , <i>pat</i> , <i>Zea mays</i> subsp. <i>mays</i> (L.) Iltis) (186165, OECD UI: DP-186165-2) |
| Content of the Type 1 Use of Living Modified Organism: | Cultivation in isolated field, storage, transportation, disposal, and acts incidental to them |
| Method of the Type 1 Use of Living Modified Organism: | <p>Location: 19-2, Kiyohara-Kogyodanchi, Utsunomiya-city, Tochigi-Ken</p> <p>Name: Isolated field, Utsunomiya Site, Du Pont Kabushiki Kaisha</p> <p>Period of Use: From March 27, 2013 to March 31, 2016</p> <p>1. Facilities of the isolated field</p> <p>(1) Security fence around the circumference of the isolated field to prevent unauthorized entry.</p> <p>(2) Prominent notice board that is easy to find, indicating the area is an isolated field and is forbiddance of unauthorized entry, also indicating the name of the administrator.</p> <p>(3) Washing area to wash out soil and recombinant maize, etc from machinery, equipment, and footwear used in the area. Install facilities in a drainage system to prevent unexpected flowing out of recombinant maize.</p> <p>(4) Prevent dispersal of seed and seedlings of recombinant maize due to feeding damage of animals such as birds by setting up bird nets from the seeding and maturing period until harvest time.</p> |

| | |
|--|---|
| | <p>2. Operation procedures in the isolated field</p> <ol style="list-style-type: none">(1) Suppress the growth of plants other than recombinant maize and the control maize to the lowest level.(2) Using secure container which prevent unexpected spillage of recombinant maize, when recombinant maize goes out from isolated field as transit and storage.(3) Inactivate recombinant maize after cultivation by plowing back in the isolated field or other methods, except transit and storage of the recombinant maize outside the isolated field in compliance with paragraph (2).(4) Prevent unexpected outspread of recombinant maize by washing machinery, equipment, and footwear used in the isolated field.(5) Implement the detasseling or staminate bagging in order to prevent pollen dispersal of recombinant maize.(6) Maintain and manage the facility so that it can adequately function.(7) Ensure personnel compliance with paragraphs (1) through (6) when implementing Type 1 Use.(8) When risk of an adverse effect on biological diversity is detected, ensure implementation of stipulated “emergency measures plan”. |
|--|---|