Corporation obtaining approval, the name of its representative, and the address of its main office

Incorporated Administrative Agency Forest Tree Breeding Center, Japan President Akira TANOOKA 3809-1 Ishi, Juochyo, Hitachi-shi, Ibaraki Prefecture

Name of the type of Living	High cellulose rich white poplar trg300-2
Modified Organism:	(AaXEG2, Populus alba L.)
Content of the Type 1 Use	Cultivation in isolated field, storage, transportation, disposal, and acts incidental
of Living Modified	to them
Organism:	
Method of the Type 1 Use	Location: 3809-1, Ishi, Juocyo, Hitachi-shi, Ibaraki Prefecture
of Living Modified	
Organism:	Name: Isolated field, Incorporated Administrative Agency
	Forest Tree Breeding Center, Japan.
	Period of Use: From March 22, 2007 to December 31, 2011
	1. Facilities of the isolated field
	(1) Security fence of 8m in height around the circumference of the isolated
	field to prevent unauthorized entry and dispersion of broken twig. The
	fence has concrete protection wall up to 1m in the underground.
	(2) Prominent notice board that is easy to find at the main entrance, indicating
	the area is an isolated field and is forbiddance of unauthorized entry, also
	indicating the name of the administrator.
	(3) Washing area to wash out recombinant white poplar from machinery,
	equipment, and footwear used in the area. Install facilities in a drainage
	system to prevent unexpected flowing out of recombinant white poplar.

Approved Type 1 Use Regulation

2. Operation procedures in the isolated field
(1) Suppress the growth of plants other than recombinant white poplar and the
control poplar to the lowest level.
(2) Using secure container which prevent unexpected spillage of recombinant
white poplar, when recombinant white poplar goes out from isolated field
as transit and storage.
(3) When fear for a part of recombinant white poplar over the fence of 8m in
height is caused, promptly cut only part of over the fence.
(4) When acknowledged recombinant white poplar make flower bud,
implement measure to prevent crossing by promptly cut theirs.
(5) Inactive recombinant white poplar and the control poplar after cultivation
by drying after cut it down, cutting and piled up at chip dump site which
made bare concrete floor inside isolated filed. After that plowing it back in
the isolated field, except transit and storage of the recombinant white
poplar outside the isolated field in compliance with paragraph (2).
Furthermore, Inactive dead branch and falling leaf by drying or burning as
work as reasonably achievable collect theirs. Inactive remain roots by
spraying glyphosate to the leaf and injection to the stump, when their root
bud cultivate until coming June after it had been cut down.
(6) Prevent unexpected outspread of extraneous recombinant white poplar by
washing machinery, equipment, and footwear used in the isolated field.
(7) Maintain and manage the facility so that it can adequately function.
(8) Ensure personnel compliance with paragraphs (1) through (7) when
implementing Type 1 Use.
(9) Implement the monitoring based on the plan of monitoring which has been
decided in addition.
(10) When risk of an adverse effect on biological diversity is detected, ensure
implementation of stipulated "emergency measures plan".