

**Corporation obtaining approval, the name of its representative, and the address of its main office**

Syngenta Seeds K. K.  
 Okifumi Murata, President  
 401-2, Mukounodai, Takatsuhara,  
 Tako-machi, Katori-gun,  
 Chiba-ken

**Approved Type 1 Use Regulation**

Name of the type of Living Modified Organism:	Maize tolerant to corn rootworm ( <i>ecry3.1Ab</i> , <i>Zea mays</i> subsp. <i>mays</i> (L.) Iltis) (Event 5307, OECD UI:SYN-Ø53Ø7-1)
Content of the Type 1 Use of Living Modified Organism:	Cultivation in isolated field, storage, transportation, disposal, and acts incidental to them
Method of the Type 1 Use of Living Modified Organism:	<p>Location: 138, Kanza, Shimada-city, Shizuoka-ken</p> <p>Name: Isolated field, Kanza site, central research station, research department, Syngenta Japan K. K.</p> <p>Period of Use: From June 11, 2010 to March 31, 2012</p> <p>1. Facilities of the isolated field</p> <p>(1) Security fence around the circumference of the isolated field to prevent unauthorized entry.</p> <p>(2) Prominent notice board that is easy to find, indicating the area is an isolated field and that unauthorized entry is forbidden; the name of the administrator is also indicated</p> <p>(3) Washing area to remove soil and recombinant maize, etc from machinery, equipment, and footwear used in the area. Install facilities in a drainage system to prevent unexpected overflow and spread of recombinant maize.</p> <p>(4) Nets surround the isolated field, functioning as a shelter against wind to reduce the dispersal of pollen.</p> <p>2. Operation procedures in the isolated field</p> <p>(1) Suppress to the lowest level the growth of plants other than recombinant and control maize.</p>

	<ul style="list-style-type: none"><li>(2) Use secure containers to prevent unexpected spillage of recombinant maize, during transportation and storage.</li><li>(3) Inactivate recombinant maize after cultivation by plowing back into the isolated field or by using other methods, except during transit and storage of the recombinant maize outside the isolated field in compliance with paragraph (2).</li><li>(4) Prevent unexpected spread of extraneous recombinant maize by washing machinery, equipment, and footwear used in the isolated field.</li><li>(5) Implement the detasseling or staminate bagging in order to prevent pollen dispersal of recombinant maize.</li><li>(6) Maintain and manage the facility so that it can adequately function.</li><li>(7) Ensure personnel comply with paragraphs (1) through (6) when implementing Type 1 Use.</li><li>(8) When risk of an adverse effect on biological diversity is detected, ensure implementation of stipulated “emergency measures plan”.</li></ul>
--	---