

**Corporation obtaining approval, the name of its representative, and the address of its main office**

Monsanto Japan Limited  
 Seiichiro Yamane, President  
 Ginza Sanno Bldg. 8F, 4-10-10  
 Ginza, Chuo-ku, Tokyo

Approved Type 1 Use Regulation

Name of the type of Living Modified Organism:	Oilseed rape tolerant to glyphosate herbicide (Modified <i>cp4 epsps</i> , <i>Brassica napus</i> L.) (MON88302, OECD UI: MON-88302-9)
Content of the Type 1 Use of Living Modified Organism:	Cultivation in isolated field, storage, transportation, disposal, and acts incidental to them
Method of the Type 1 Use of Living Modified Organism:	<p>Location: 4717, Manaita, Kawachi-machi, Inashiki-gun, Ibaraki-Ken</p> <p>Name: Isolated field, Monsanto Japan Limited</p> <p>Period of Use: From August 8, 2011 to May 31, 2014</p> <p>1. Facilities of the isolated field</p> <p>(1) Security fence around the circumference of the isolated field to prevent unauthorized entry.</p> <p>(2) Prominent notice board that is easy to find, indicating the area is an isolated field and is forbiddance of unauthorized entry, also indicating the name of the administrator.</p> <p>(3) Washing area to wash out soil and recombinant oilseed rape, etc from machinery, equipment, and footwear used in the area. Install facilities in a drainage system to prevent unexpected flowing out of recombinant oilseed rape.</p> <p>(4) Nets are surrounding the isolated field, functioning as a shelter against wind to reduce the dispersal of pollen. Setting up cheesecloth at the flowering</p>

period to decrease the pollen dispersal.

- (5) When sowing season and maturing period, implement counter measure that using bird nets, etc to prevent bird injury.

2. Operation procedures in the isolated field

- (1) Suppress the growth of plants other than recombinant oilseed rape and the control oilseed rape to the lowest level.
- (2) Using secure container which prevent unexpected spillage of recombinant oilseed rape, when recombinant oilseed rape goes out from isolated field as transit and storage.
- (3) Inactivate recombinant oilseed rape after cultivation by plowing back in the isolated field or other methods, except transit and storage of the recombinant oilseed rape outside the isolated field in compliance with paragraph (2).
- (4) Prevent unexpected outspread of extraneous recombinant oilseed rape by washing machinery, equipment, and footwear used in the isolated field.
- (5) Maintain and manage the facility so that it can adequately function.
- (6) Ensure personnel compliance with paragraphs (1) through (5) when implementing Type 1 Use.
- (7) Implement the monitoring based on the plan of monitoring which has been decided in addition.
- (8) When risk of an adverse effect on biological diversity is detected, ensure implementation of stipulated “emergency measures plan”.