## Corporation obtaining approval, the name of its representative, and the address of its main office

Du Pont Kabushiki Kaisha Minoru Amoh, President 2-11-1, Nagata-chou, Chiyoda-ku, Tokyo

Approved Type T Use Keg	
Name of the type of Living	Maize resistant to Lepidoptera and Coleoptera and tolerant to
Modified Organism:	glufosinate herbicide
	(Modified cry1F, cry34Ab1, cry35Ab1, pat, Zea mays subsp. mays (L.) Iltis)
	(43A47, OECD UI: DP-Ø43A47-3)
Content of the Type 1 Use	Cultivation in isolated field, storage, transportation, disposal, and acts incidental
of Living Modified	to them
Organism:	
Method of the Type 1 Use	Location: 19-2, Kiyohara-Kogyodanchi, Utsunomiya-city, Tochigi-Ken
of Living Modified	
Organism:	Name: Isolated field, Utsunomiya Site, Du Pont Kabushiki Kaisha
	Period of Use: From September 2, 2011 to March 31, 2014
	1. Facilities of the isolated field
	(1) Security fence around the circumference of the isolated field to prevent
	unauthorized entry.
	(2) Prominent notice board that is easy to find, indicating the area is an isolated
	field and is forbiddance of unauthorized entry, also indicating the name of
	the administrator.
	(3) Washing area to wash out soil and recombinant maize, etc from machinery,
	equipment, and footwear used in the area. Install facilities in a drainage
	system to prevent unexpected flowing out of recombinant maize.
	(4) Prevent dispersal of seed and seedlings of recombinant maize due to
	feeding damage of animals such as birds by setting up bird nets at seeding
	and maturing period.

Approved Type 1 Use Regulation

2.	Operation procedures in the isolated field
(	1) Suppress the growth of plants other than recombinant maize and the control
	maize to the lowest level.
(2	2) Using secure container which prevent unexpected spillage of recombinant
	maize, when recombinant maize goes out from isolated field as transit and
	storage.
(.	3) Inactivate recombinant maize after cultivation by plowing back in the
	isolated field or other methods, except transit and storage of the
	recombinant maize outside the isolated field in compliance with paragraph
	(2).
(4	4) Prevent unexpected outspread of recombinant maize by washing machinery,
	equipment, and footwear used in the isolated field.
	5) Implement the detasseling or staminate bagging in order to prevent pollen
	dispersal of recombinant maize.
((	6) Maintain and manage the facility so that it can adequately function.
(	7) Ensure personnel compliance with paragraphs (1) through (6) when
	implementing Type 1 Use.
3)	3) When risk of an adverse effect on biological diversity is detected, ensure
	implementation of stipulated "emergency measures plan".