

Corporation obtaining approval, the name of its representative, and the address of its main office

Monsanto Japan Limited
 Seiichiro Yamane, President
 Ginza Sanno Bldg. 8F
 4-10-10, Ginza, Chuo-ku, Tokyo

Approved Type 1 Use Regulation

Name of the type of Living Modified Organism:	Sugar beet tolerant to glyphosate herbicide (<i>cp4 epsps, Beta vulgaris L. ssp. vulgaris var altissima</i>) (H7-1, OECD UI : KM-000H71-4)
Content of the Type 1 Use of Living Modified Organism:	Cultivation, storage, transportation, disposal and acts incidental to them, in isolated fields.
Method of the Type 1 Use of Living Modified Organism:	<p>Location: 4475-2, Aza-Dotemukai, Manaita, Kawachi-machi, Inashiki-gun, Ibaraki-ken</p> <p>Name: Isolated field, Monsanto Japan Limited</p> <p>Period of use: From May 25, 2005 to December 31, 2005</p> <p>1. Facilities of the isolated field</p> <p>(1) Security fence around the circumference of the isolated field to prevent unauthorized entry.</p> <p>(2) Prominent notice board that is easy to find, indicating the area is an isolated field and is forbiddance of unauthorized entry, also indicating the name of the administrator.</p> <p>(3) Washing area to wash out living modified organisms from machinery, equipment, and footwear used in the area. Install sedimentation tank and nets in a drainage system to prevent unexpected flowing out of extraneous recombinant Sugar beet.</p>

	<p>(4) Installation of a windbreak net around the circumference of the isolated field.</p> <p>2. Operation procedures in the isolated field</p> <p>(1) Suppress the growth of plants other than recombinant Sugar beet and the control Sugar beet to the lowest level.</p> <p>(2) Sow seeds and grow seedlings in a greenhouse at the Kawachi Research Farm of Monsanto Japan Limited. Transplant the seedlings to the isolated field at 2 to 4 leaf stage. Sow one seed per pod, and dispose the seeds, which do not germinate, by plowing into the isolated field, after autoclaving the pods in order to inactivate those seeds completely. After the completion of cultivation of recombinant plants, inactivate by plow-in or incineration in incinerator.</p> <p>(3) Prevent unexpected spillage of recombinant Sugar beet in transit and storage by using secure container.</p> <p>(4) Inactivate recombinant Sugar beet after cultivation by plowing back in the isolated field or other methods, except transit and storage of the recombinant Sugar beet outside the isolated field in compliance with paragraph (3).</p> <p>(5) Prevent unexpected outspread of extraneous recombinant Sugar beet to machinery, equipment, and footwear used in the area by washing out in the isolated field.</p> <p>(6) Maintain and manage the facility so that it can adequately function.</p> <p>(7) Ensure personnel compliance with paragraphs (1) through (6) when implementing Type 1 Use.</p> <p>(8) When risk of an adverse effect on biological diversity is detected, ensure implementation of stipulated “emergency measures plan”.</p>
--	--